

# Archaeo 2

---

The Chronostratigraphic Record  
and Methods of Relative and  
Absolute Dating in Archaeology

# Time line of the genus *homo*



Homo floresiensis

## Genealogical Tree of Hominids





Abbildung 2: Stammbaumphypothese zur Evolution des Menschen auf biogeographischer Grundlage.


"Hominid family". Left to right: Top row: Kenyanthropus platyops, Homo neanderthalensis; middle row: Australopithecus afarensis, Paranthropus boisei, Homo habilis; bottom row: Australopithecus africanus, Homo erectus, Australopithecus anamensis, Homo rudolfensis.



# Time line of the genus *homo*

## Genealogical Tree of Hominids

A. = *Australopithecus*  
H. = *Homo*



# Stratigraphy - Nicholas Steno (1638-1686)

Pioneered in Anatomy and Geology


Principles of Stratigraphy (1669)

1. Superposition
2. Original horizontality
3. Lateral continuity and cross-cutting discontinuities





Illustration from Steno's 1667 paper comparing the teeth of a shark head with a fossil tooth

# Stratigraphy - Nicholas Steno




ogy

569)



## One more stratigraphic principle

---



Principle of palaeontological identity: Two layers with the same fossil contents have the same age.

---

# Chronology of Geology and Naturalism

---

## Charles Lyell (1797-1875)

Principles of Geology (1830):

Continuity and uniformitarianism - the Earth was shaped by the same processes that are still in operation today




## Charles Darwin (1809-1882)

Origin of Species (1859):

Evolution

Faunal assemblages change over time



# Historical Dating

---



Roman coin dating to the reign of Nero 54-68 AD,

The recovery of material of a known age from a site, i.e. coins, bottles, ceramics, beads can be used to date the site itself.




Celadon bowl, Yuan/Ming Dynasty China, 14<sup>th</sup> century AD

---

# Historical Dating

---




TPQ- *Terminus post quem* or the earliest possible date for an archaeological deposit

TAQ- *Terminus ante quem* or the latest possible date for the deposit

# Laguna Copper Plate

---




Legal document written in Old Javanese/Old Tagalog script

The document bears a date of Saka 844 or **922 AD**

---

# Prehistoric Chronology

---




The stratigraphic method

---

# Relative Chronology

---



**Oskar Montelius (1901)**

“Relative Chronology tells us if an object is younger or older than another object.”

---

# Relative Chronology

Historic Accounts in combination with Archaeology:

Predynastic and early Dynastic Egyptian Kings and Tombs

date BC	period	(H.) phase	(K.) phase	king	Lower Nubia	Hierakonpolis	Naqada	Abadiya	Abydos	Lower Egypt
3550	Naqada I	IC-IIA	Ic-IIa			Locality 6	1587		Cemetery U	
						Tomb 3 Tomb 6	1503 1497 1610	Cemetery B B101 B102	U-239	
3500	Naqada II	IIB	Iib				1426			
3450							1411			
3400			IIC	Iic			Painted Tomb T100	T4 Cemetery T		
3350							Cemetery	T5 T3 T32		
3300		IID1	IId1				T9 T10 T16 T17		U-q	
3250		IID2	IId2-IIIa1				T25		U-547	
3200	Naqada III	IIIA1	IIIA2							
3150			IIIA2			Qustul Cem L, Sevata L24 137.1		Gebel Tjauti		U-j
3100		'Dyna 00' 'Dyna 0'	IIIB	IIIB1			Gebel Sheikh Suleiman	Wadi Qash Site 34		U-s
3050	First Dynasty		IIIB2						B7/9	
3000			IIIC1	IIIC1	A X B Ka Y Nar					
2950				IIIC2	Aha Djer		Narmer palette	royal tombs	B17/18	Buto sealing

T.A.H. Wilkinson (M.D.A.I.K. 56, 2000 p. 392)

Fig. 5: Summary chart of political unification

Dates BC are approximate. The two columns headed 'phase' give the divisions of the Predynastic cultural sequence devised by HENDRICKX (1996) and KAISER (1957, 1990), respectively

# Relative Chronology

Different Kings or Different Names?

Historic Accounts in combination with Archaeology:

They can be the same person. Predynastic and early Dynastic

Egyptian Kings and Tombs with Pharaoh Menes, the legendary founder of the First Dynasty

However, it is not always a straightforward process, and often hampered by the ambiguity of records and different interpretations

The Scorpion King could have been an unknown King who ruled in Egypt between the reigns of Ka and Narmer


date BC	period	(H.) phase	(K.) phase	king	Lower Nubia	Hierakonpolis	Naqada	Abadiya	Abydos	Lower Egypt
3550	Naqada I	IC-IIA	IC-IIa			Locality 6 Tomb 3 Tomb 6	1587		Cemetery B B101 B102	Cemetery U U-239
3500							1426			
3450	Naqada I	IIB	IIb							
3400										
3350										
3300										
3250										
3200										
3150										
3100										
3050										
3000										
2950										



wan  
khan  
ra  
  
to  
aling  
  
ed by


# Relative Chronology

---



The Tomb of King Narmer in Umm-al-Qaab, Abydos

Map of pharaonic tombs at Abydos



# Relative Chronology

---



The Scorpion King Macehead



Palette of King Narmer

# Relative Chronology

---

Known rulers in the history of Egypt for the First Dynasty are as follows:

Name	Comments	Dates
Narmer	- probably Menes on earlier lists	c. 3100–3050 B.C.
Hor-Aha		c. 3050–3049 B.C.
Djer	-	c. 3049–3008 B.C. 41 years (Palermo Stone)
Djet	-	3008–2975?
Merneith	the mother of Den	3008?
Den	-	2975–2935 30 to 50 years (40 years?)
Anedjib	-	2935?–2925? 10 years (Palermo Stone)
Semerkhet	-	2925?–2916? 9 years (Palermo Stone)
Qa'a	-	2916?–2890 B.C.

---

# Relative Chronology

---

Chronology of Technological Developments

E.g. Pyramids



# Relative Chronology


---




Mastaba of Ptahchepses



Mastaba at el-Faraoun



Reconstruction of a 1st Dynasty Mastaba



Typical architectural design of a Mastaba

# Relative Chronology



---



---

Step Pyramid of King Djoser in Sakkara



# Relative Chronology



The "Pyramid King": 5-Step Pyramid of Pharaoh Sneferu in Meidum

# Relative Chronology

---



The "Pyramid King": The "Bent Pyramid"

# Relative Chronology

---



---

The "Pyramid King": The "Red Pyramid" of Pharaoh Sneferu in Dashur

# Relative Chronology

---




---

Finally: The Great Pyramid of Khufu in Giza

# Relative Dating Methods

---

## □ Typology



Typological Method:


Tells us whether an object is older or younger than another object

Typological Rudiment:

If an element of an object that originally had a practical function loses it, but lives on as an ornament, then it is a typological rudiment

---

# Relative Dating Methods



**Typological Rudiment:**  
If an element of an object that originally had a practical function loses it, but lives on as an ornament, then it is a typological rudiment




# Typology

---

Egyptian pottery:

Changes and evolution of  
pottery styles and decors

After Flinders Petrie



# Typology






---

Southwest American Hohokam pottery:

Bowl styles - A 500 year sequence of Pottery typology based on painted decoration


Relative Chronology through  
**Seriation**

---

PHASE	DECORATION	SHAPE
SACATON AD 1000-1175		
SANTA CRUZ AD 875-1000		
GILA BUTTE AD 800-875		
SNAKETOWN AD 750-800		
SWEETWATER AD 700-750		
ESTRELLA AD 650-700		

# Seriation - Chart for Pottery Types


---



If one finds a lot of Beakers and few Grooved ware ,  
the site can be relatively dated to c. 2500 BC



# Seriation - Chart for Pottery Types

---




If one finds a lot of Peterborough ware and few Grooved ware, the site can be relatively dated to c. 3000 BC

# Seriation - Chart for Pottery Types



of Grooved ware  
 ted to c. 2700 BC  
 Computer software WinSerion for Seriation






## What we date

Carbonates

Teeth and bones of  
Big mammals

Sediments



Volcanic minerals

Wood and charcoals



Burned elements

# Absolute Dating

---

- Palaeomagnetism
  - Dendrochronology
  - Radiocarbon Dating
  - Electron-Spin-Resonance
  - Potassium-Argon dating
  - Thermoluminescence / Optoluminescence
  - Uranium Series
-

# Palaeomagnetism




# Dendrochronology

---

A cross-section of a downed tree, showing annual growth rings.




# Dendrochronology



**Figure 4.9** The key principles of dendrochronology

Principle of tree ring dating: overlaps are matched to take the sequence back from a known date to date old timbers. Samples are taken at 90 degrees to the grain and numbers of rings and their thickness measured by eye or computer.

# Dendrochronology



# Dendrochronology

---



# Radiocarbon Dating


All living organisms absorb Carbon and its isotopes from the atmosphere - until they die.

Radioactive Carbon 14 ( $^{14}\text{C}$ ) decays at a known rate or half-life (5730 years). Calculation of the amount of the remaining  $^{14}\text{C}$  provides a date for dead matter


The level of cosmic radiation has fluctuated over time, hence the need to calibrate.

Calibration routines used to convert radiocarbon years to calendar years  
A statistical estimation of error is expressed as standard deviation

Sample size for conventional  $^{14}\text{C}$  dating is 10-20g, for Accelerator Mass Spectrometry (AMS) is 0.1g



# Radiocarbon Dating



OxCal 4.2 Manual

File View Format Options Data Help

Name

Date

±

Curve IntCal 13

Calibrate

View Curve

Projects...

Interface build: 78 Updated: 9/10/2013  
(c) Christopher Bronk Ramsey 2013

OxCal 4.2 Manual

- Introduction
- Licence
- Installation
- Input of information
- Analysis
- Calibration Curves
- Viewing output
- Command reference
- Program development history
- References and bibliography
- Acknowledgements

Web interface build number: 78  
Last Updated: 9/10/2013  
(c) Christopher Bronk Ramsey 2013

References

- Bronk Ramsey 1994 (first release notice)
- Bronk Ramsey 1995 (main introduction to program)
- Bronk Ramsey 1998 (approach and plans)
- Bronk Ramsey 2001 (developments 1995-2001)
- Bronk Ramsey et al. 2001 (D\_Sequence method and tests of other methods)
- Bronk Ramsey 2008 (deposition models) [pre-print]
- Bronk Ramsey 2009a (new features of OxCal4)
- Bronk Ramsey 2009b (outlier analysis) [pre-print]
- Bronk Ramsey et al. 2010
- Lee and Bronk Ramsey 2012 (trapezoidal models)
- Radiocarbon Vol 55(4) (IntCal13 special issue)

If you use this program, you should quote the reference for the calibration curve used, the version of OxCal (with any non-standard options set) and the reference Bronk Ramsey 2009a. If you are wiggle-matching tree-ring sequences you should quote Bronk Ramsey et al. 2001. For deposition models Bronk Ramsey 2008 should be referenced and for outlier analysis Bronk Ramsey 2009b. For use of trapezoidal priors please quote Lee and Bronk Ramsey 2012.

**Introduction**

OxCal is a program designed for the analysis of chronological information.

**Radiocarbon**  
An International Journal of Cosmogenic Isotope Research  
VOLUME 55 / NUMBER 4 / 2013  
14  
INTCAL13  
P. J. REIMER, GUEST EDITOR

<https://c14.arch.ox.ac.uk/oxcal/OxCal.html>

# Radiocarbon Dating



Radiocarbon Laboratory  
Australian National University

*The University of Waikato*  
*Radiocarbon Dating Laboratory*




Private Bag 3105  
Hamilton,  
New Zealand.  
Fax +64 7 838 4192  
Ph +64 7 838 4278  
email c14@waikato.ac.nz  
Head: Dr Alan Hogg

## Report on Radiocarbon Age Determination for Wk- 32984

<b>Submitter</b>	P Piper
<b>Submitter's Code</b>	IV-2011-G3-573
<b>Site &amp; Location</b>	Bubog Island, Philippines
<b>Sample Material</b>	Conus sp.
<b>Physical Pretreatment</b>	Surfaces cleaned. Washed in an ultrasonic bath. Tested for recrystallization: aragonite.
<b>Chemical Pretreatment</b>	Sample acid washed using 2 M dil. HCl for 120 seconds, rinsed and dried.

$\delta^{13}\text{C}$	0.2 ± 0.2 ‰
$\text{D}^{14}\text{C}$	-483.4 ± 2.5 ‰
$\text{F}^{14}\text{C}\%$	51.7 ± 0.2 ‰
<b>Result</b>	<b>5306 ± 38 BP</b>

### Comments



**Result after calibration**

*Alan Hogg*  
21/02/12

- Result is *Conventional Age or Percent Modern Carbon (pMC)* following Stuiver and Polach, 1977, Radiocarbon 19, 355-363. This is based on the Libby half-life of 5568 yr with correction for isotopic fractionation applied. This age is normally quoted in publications and must include the appropriate error term and Wk number.
- Quoted errors are 1 standard deviation due to counting statistics multiplied by an experimentally determined Laboratory Error Multiplier.
- The isotopic fractionation,  $\delta^{13}\text{C}$ , is expressed as ‰ wrt PDB.
- $\text{F}^{14}\text{C}\%$  is also known as *Percent Modern Carbon (pMC)*

# Radiocarbon Dating



Radiocarbon Laboratory  
Waikato University

*The University of Waikato*  
*Radiocarbon Dating Laboratory*




Private Bag 3105  
Hamilton,  
New Zealand.  
Fax +64 7 838 4192  
Ph +64 7 838 4278  
email c14@waikato.ac.nz  
Head: Dr Alan Hogg

## Report on Radiocarbon Age Determination for Wk- 32984

<b>Submitter</b>	P Piper
<b>Submitter's Code</b>	IV-2011-G3-573
<b>Site &amp; Location</b>	Bubog Island, Philippines
<b>Sample Material</b>	Conus sp.
<b>Physical Pretreatment</b>	Surfaces cleaned. Washed in an ultrasonic bath. Tested for recrystallization: aragonite.
<b>Chemical Pretreatment</b>	Sample acid washed using 2 M dil. HCl for 120 seconds, rinsed and dried.

$\delta^{13}\text{C}$	$0.2 \pm 0.2 \text{‰}$
$\text{D}^{14}\text{C}$	$-483.4 \pm 2.5 \text{‰}$
$\text{F}^{14}\text{C}\%$	$51.7 \pm 0.2 \%$
<b>Result</b>	<b>5306 <math>\pm</math> 38 BP</b>

### Comments



**Result after calibration**

*Alan Hogg*  
21/02/12

- Result is *Conventional Age* or *Percent Modern Carbon (pMC)* following Stuiver and Polach, 1977, Radiocarbon 19, 355-363. This is based on the Libby half-life of 5568 yr with correction for isotopic fractionation applied. This age is normally quoted in publications and must include the appropriate error term and Wk number.
- Quoted errors are 1 standard deviation due to counting statistics multiplied by an experimentally determined Laboratory Error Multiplier.
- The isotopic fractionation,  $\delta^{13}\text{C}$ , is expressed as ‰ wrt PDB.
- $\text{F}^{14}\text{C}\%$  is also known as *Percent Modern Carbon (pMC)*

# Radiocarbon Dating



# Electron Spin Resonance (ESR)


---

- ❑ Electric charges build up at known rate in some crystal structures. The time since the process began can be calculated by measuring the charge produced when subjected to microwave energy.
  - ❑ Range: 50 ky to 1my
  - ❑ Teeth enamel, shells and calcite deposits in caves
-

# Thermoluminescence (TL)

## Optical Stimulated Luminescence (OSL)

Radioactive decay in quartz crystals leads to a build up of electric charges at a known rate. This electrical charge is released as light when the crystals are heated for TL or light induced for OSL.



The amount of energy released is relative to the amount of time since last heated to over 500 °C the 'clock setting event'.



## Espinosa Fossil Site, Cagayan Valley

---

OSL sample collection

---






Gamma ray mass spectrometry  
of the sediment  
Estimating background  
radiation



# Thermoluminescence (TL) / Optical Stimulated Luminescence (OSL)













# Potassium-Argon ( $^{40}\text{K}/^{40}\text{Ar}$ )

## Argon-Argon ( $^{40}\text{Ar}/^{39}\text{Ar}$ )

---

Potassium decays to its daughter element Argon with a half life of 1.25 ma.

Argon-Argon requires neutron activation in a research reactor. It has a precision advantage compared to  $^{40}\text{K}/^{40}\text{Ar}$  for younger periods (100ky - 10ky)

Applicable for dating volcanic rocks such as ignimbrites containing hornblende and sanidine

---



# Uranium Series Dating

---



Using uranium isotopes marks the beginning of radiometric dating. Arthur Holmes developed the uranium-lead dating method already in 1911 at the age of 21, one year after graduation.

The U-Pb method relies on two separate decay chains, the uranium series from  $^{238}\text{U}$  to  $^{206}\text{Pb}$ , with a half-life of 4.47 billion years and the actinium series from  $^{235}\text{U}$  to  $^{207}\text{Pb}$ , with a half-life of 704 million years.

It can be used to date rocks of an age between 1ma to over 4500ma. Applied for the determination of the age of the earth. Mostly not applicable in Archaeology.

---

# Uranium-Thorium Dating

---

Uranium-234 is water-soluble and occurs in all natural waters. It decays to Thorium-230 which is not water-soluble with a half-life of 245ky.

Thorium-230 is itself radioactive with a half-life of 75ky and so instead of accumulating indefinitely it approaches a secular equilibrium with its parent isotope.

At equilibrium, the number of Thorium-230 decays per year within a sample is equal to the number of Uranium-234 decays per year in the same sample.

Can be used to date calcium carbonate material, i.e. cave speleothem, corals, fossilized bones and teeth

Upper age limit is 500ky

---

# Uranium-Thorium Dating

U/Th Dating of human fossils from Tabon Cave: Low collagen contents prevented  $^{14}\text{C}$  dating. U/Th method permitted a direct dating of the human bones. Redated „Tabon Man“ from c. 22ky to 16.5ky

Sample	U (ppm)	$^{234}\text{U}/^{238}\text{U}$	$^{230}\text{Th}/^{232}\text{Th}$	$^{230}\text{Th}/^{234}\text{U}$	Age (Kyr BP)
Frontal bone P-XIII-T-288	2.88	$1.115 \pm 0.069$	> 100	$0.142 \pm 0.016$	$16.5 \pm 2.0$
Right mandibular fragment PXIIIT436-Sg19	0.56	$1.169 \pm 0.210$	48	$0.249 \pm 0.049$	$31 +8/-7$
Tibia fragment (IV-2000-T-197)	1.11	$1.174 \pm 0.150$	53	$0.354 \pm 0.061$	$47 +11/-10$



courtesy C. Falguères (IPH, Paris)

# Geochronological Dating

---



# Geochronological Dating

---

## □ Loess

### ■ Time capsules in the dust

Aeolian deposits:

Smallest particles of silt and sand are picked up from exposed surfaces.

Vegetation-free polar deserts of high latitudes and alpine mountain ranges are especially suited to wind abrasion.

- Grain size between 10-50 $\mu$ m
- Non stratified, homogenous deposit with porous structure
- Composition dominated by quartz grains
- Originates in periglacial context (cold arid to semi-arid, i.e. cold steppe)
- Loessification: process of syndepositional diagenesis



Sand storm in the Gobi desert (Mongolia), June 2008

# Geochronological Dating

## □ Loess

- Time capsules in the dust: Palaeosols captured by aeolian deposits




Loess bluff at Edwardsville, Illinois, USA



Eemian ~128-117ky

Bucket excavator in Lower Rhine Basin, Germany




Loess landscape in Shanxi, China

# Geochronological Dating

## □ MIS/OIS

### ■ Time capsules in marine sediments and ice

- Loess deposition is related to periglacial environments during glacial maxima (pleniglacial stages),
- Geological features and structures related to cold climate and permafrost (topsoil is frozen during the entire year and thus watertight)
- Record of climatic variations. During climatic ameliorations, vegetation growth leading to soil formation. These soils are characteristic for certain climatic and biological environments (e.g. Eemian)
- Oxygen Isotope Stages are cyclical variations in the ratio of the abundance of oxygen with an atomic mass of 18 to the abundance of oxygen with an atomic mass of 16. The ratio is linked to water temperature of ancient oceans, which in turn reflects ancient climates.



# Combining Relative and Absolute Dating


---

## □ Interdisciplinary Approach

Archaeology in correlation with environmental and geological records:

- Typology
  - Stratigraphies
  - Pollen analysis
  - Archaeozoology
  - OIS/MIS sequences
-





Correlation of geological, pedological and biological stratigraphic sequences for the Upper Pleistocene

# Group work

---

- Groups of 3-4 people
  - Review the methods of Relative and Absolute Dating
  - Submit a report
-